RECEIVED
FEB 2 0 2007
GEOSCIENCE ASSESSMENT
OFFICE

2.34229

Plinky Property

West Group

Eby Township

NTS - 42 A/1

80°10'W 48°04'N

EDH-12

Prepared by : Eric Marion

INDEX

Property Location.....3

Access 3

Claims.....3

General Geology.....3

Claim Geology.....5

Previous Work.....5

Present Work.....9

Result/Conclusions.....9

Drill Hole Log.....10

Assay Sheets.....appended

Bibliography.....i , ii

FIGURES

Fig - 1 Location.....1

Fig - 2 Claim Map.....2

Fig - 3 General Geology.....4

Fig - 4 Claim Geology.....6

Fig - 5 Drill Hole Location.....7

Fig - 6 Drill Hole Section EDH -12.....8

KIRKLAND LAKE RESIDENT GEOLOGIST'S'S DISTRICT




Figure 1




Figure - 2

Eby Twp. Plan M-0345

2

UTM Zone 17
1000m grid

PROPERTY LOCATION

The property is located in the Larder Lake mining division approximately 12 kilometers south-west of the town of Kirkland Lake. The group sits midway in the eastern end of the township.This is in the Kirkland Lake Resident Geologists District and can be found on NTS-42 A/1 with the geographic center being at approximately 80°10'W and 48°04'N.

ACCESS

Heading south on Hwy.#11 from the intersection of Hwy.#66 and trans-Canada Hwy.#11 at Kenogami for 4 kilomneters will bring you to the approximate geographic center of the group. Many hunting trails, transmission lines and a gas pipeline give improved access to much of the claim area.


CLAIMS

This property consists of eight contiguous, staked mining claim blocks totalling 28 claim units in the subdivided township of Eby, recorded on Plan M-0345 of Eby Twp.The claims and descriptions are as follows:

CL#1239348--Lot 3, Con.5 N1/2(noSW1/4) +Lot 3 Con.5 N1/2 of S1/2
+Lot 2, Con.5 NW1/4 of S1/2 - 6 units
CL#1239349--Lot 2, Con.5 S1/2 of S1/2 - 2 units
CL#1239350--Lot 1, Con.4 NW1/4 of N1/2 - 1 unit
CL#1239352--Lot 3, Con.4 N1/2(noSW1/4) + Lot 2, Con.4 SW1/4 of N1/2 - 4 units
CL#1239353--Lot 1, Con.4 SW1/4 of N1/2 = Lot 2,Con.4 SE1/4 of N1/2
+ Lot 2, Con.4 NE1/4 of S1/2 - 3 units
CL#1239354--Lot 3, Con.4 S1/2 + Lot 2, Con.4 W1/2 of S1/2 - 6 units
CL#3010057--Lot 2, Con.4 N1/2 of N1/2 - 2 units
CL#4202851--Lot 2, Con.3 N1/2 - 4 units

GENERAL GEOLOGY

This area lies within the Abitibi Greenstone Belt, a region of predominantly volcanic rocks and related interflow sediments at the south central region of the Superior Province.Several eras of intrusion and deformation have affected most of the lithologies present.Major structural deformation zones,(locally the Larder-Cadillac Deformation Zone), parallel each other west to east across the belt and seem to have acted as a control on gold deposition.The Abitibi Belt is host to many large gold and base metal deposits on both sides of the Ontario-Quebec border along these structural trends and has an exploration history going back well into the 1800's.A band of altered mainly fluvial sediments of Temiskaming age, folded and upturned to a near vertical position, coincide with the main structural trend of the LCDZ less than 800 meters to the north of the claim group.The round Lake batholith, a large Archaen aged granitic intrusive occurs about 2500 meters to the south.The south east extremity of the claim group touches on the "Otto Stock", an almost circular, somewhat zoned mafic(sanukitoid) intrusive of some 10 kilometer diameter.Thin bands of clastic sediments and iron formation belonging to the older Skead group trend east-west through the map area and wrap around the Otto Stock.To the west of the claim group, a roughly 10 kilometer wide north-south finger of Huronian aged sediments filling a paleo depression of probable structural origin overlie the volcanics.Field work by the OGS has shown LCDZ strain and faulting affecting these much younger overlying sediments.The Temiskaming Rift is a regional graben feature strinking at about 330° across this part of Ontario.This young rift system has a definite control association with diamond bearing intrusives such as kimberlites. Several NNW-SSE trending fault features passing through the area have been identified as probable Temiskaming Rift associated features.



LEGEND

Proterozoic

Keeweenawan diabase (not shown)

12 Cobalt Group

Archean

Matachewan diabase (not shown)

Granitic rocks

11 Granodiorite, monzonite, quartz monzonite, syenite

10 Massive to gneissic quartz diorite, tonalite, trondhjemite

Upper Supergroup

9 9a* Timiskaming Group, 9b** Destor-Porcupine Complex

8 8a, 8n, Blake River Group, 8c*** Blake River (Upper Fm., Tisdale Group)

7 7a, 7b, Kinojevis Group, 7c Kinojevis Group, (Middle Fm., Tisdale Group)

6 6a Larder Lake Group, 6b Stoughton Roquemaure Group, 6c Lower Fm., Tisdale Group

5 5c Porcupine Group

Lower Supergroups

4 4a Skead Group, 4b Hunter Mine Group, 4c Upper Fm., Deloro Group

3 3a Catherine Group, 3c Middle Fm., Deloro Group

2 2a Wabewawa Group, 2c Lower Fm. Deloro Group

1 1a Pacaud tuffs****

*a refers to Kirkland Lake Area, south limb of synclinorium (Jensen 1978c, 1979).
**b refers to Kirkland Lake Area, north limb of synclinorium (Jensen 1976, 1978b).
***c refers to Timmins Area (Pyke, 1980).
**** (Goodwin, 1965).

Figure-3 - General Geology

CLAIM GEOLOGY

The claim group lies along the north western contact of the Otto Stock, a syenitic intrusive complex which displays some large scale zoning features. The stock is almost circular with a diameter of about 12 kilometers. The majority of the rock north of the stock, would be mafic to intermediate tholeiitic volcanic and interflow sediments with lesser komatiitic flows, previously mapped as Skead Group volcanics. A band of felsic tuffs and agglomerates and some interlayered chemical sediments and iron formation parallel the contact of the stock and cut across the claim group in an arc. On the southern end of the claims, #3 post region of CL# 4202851 in an old drill hole and an unsampled 90 meter wide band of amphibolite gneiss (plus some altered gabbro) well mineralized with magnetite, pyrrhotite and odd chalcopyrite is logged as occuring between the sediments and the stock. Recent work by the OGS and the resident geologist office 1700 meters to the norteast of the noted drill hole, along highway #11, has documented anomolous gold, platinum, palladium and copper values in an amphibolite rock that may be the northeastern extension of the zone intersected in the drill hole. This feature does not outcrop in the region of the drill hole or along strike on the claims. The western end of the claim group (most of CL# 1239354) cover a previously mapped gabbroic intrusive which exibites layering or zoning of fine and courser grain phases. Sampling aroung the west end of the gabbro by a previous holder (Butte Canyon Resources) found elevated platinum/palladium values. Some epidote associated with quartz vains and some minor pyrites were noted on an outcrop at what would be the northern end of this unit.

Several faults which parallel the Larder Cadillac Defermation Zone (LCDZ) cross the claim group. A fault at the north end of the group has been defined as splay of the LCDZ and locally desplays defermation, quartz vaining and intense carbonate alteration. Gold and copper anomolies occur along this structural trend.

Flow tops and pillow facings in the area generally face to the north west on the claim group. Many small syenitic bodies outcrop in the area, some with considerable pyrite mineralization. Generally, much chloritization, carbonatizaion, sulphidization, and quartz veining occurs throughout the volcanic rocks on the claim group. Historic records note gold values on different workings of the area, but field work by the author has failed to positively locate these workings and duplicate the previous assays.

PREVIOUS WORK

Core drilling, geophysical surveys, trenching, mapping, sampling and geological studies have been conducted in the search for gold in areas on and around the claims.A fair amount of prospecting has been done on these claims as attested to by the number of unrecorded trenches and pits on quartz stringers and pyrite concentrations scattered across the property.There is quite a sizable sum of exploration work submitted as assesment work for the area on, and surrounding these claims.A mapped syncline crosses the south end of the claim group across CL# 1239354. Stripping work submitted by prior holders along this fold has exposed metasediments with bands of quartz pebble conglomerate quite similar in age and composition to "Witwatersrand Type" quartz pebble conglomerate. Diamond drilling by the owners of the claim immediately to the east of CL# 1239350 also encountered this band of sediments with some elevated copper and zinc values. The fold axis would cut diagonally across CL# 1239254 from the #1 post to the #3 post. Previous geophysical coverage of a large area of these claims have defined several anomolies along this fold axis. None of these anomolies have been drill tested. To the north of the present drill location, four short drill holes scattered on an east-west line were put in by previous claim holders.Anomolous zinc and gold were recorded in two of the holes. As claim holder, the author has resampled many workings, performed stripping and trenching as well as drilling 12 XRay size core holes (including this one) of depths of up to 120 feet to test different targets mostly defined by previous operators ground based geophysical surveys.

(1)	1930-48	Todora Kirkland/ Macassa	geol.. 11 ddhs, 121 ft. shaft	Cpy, Mo. Au in qtz.. up to 13.39% Cu
(2)	1938	A. Johnson	3 pits	up to 0.1 to 0.3 oz. Au/ton
(3)	1945	Sylvanite	trenches	trace to 0.023 oz. Au/ton
(4)	1955-57	Mr. Ahola	2 shafts, trenches, 1 ddh	up to 0.2 oz. Au/ton in tuff and qtz. veins
(5)	1965-69	M. Fishkin	3 ddhs	cpy + abund. py, cpy in south hole
(6)	1965-66	Keevil	geol.. mag, EM, 3 ddhs	8 conductors, iron fm with 15%-60% sulphides 0.005 oz. Au/ton in sludge
(7)	1978-80	P. Harrington	5 ddhs	-
(8)	1980	KLIP	1 OVB hole	3,200 ppb Au in basal till
(9)	1980	Noranda	mag and EM, 1 ddh	3 short, strong conductors, up to 0.016 oz. Au/ton over 1 m
(10)	1980-84	J. Reed	geol.. mag, EM, 3 ddhs	0.005 oz. Au/ton in grab, up to 0.01 oz. Au/ton over 1.5 m in core
(11)	1984-86	P. Rivard	trenching, 2 ddhs	up to 0.04 oz. Au/ton in grabs
(12)	1971	T GUTARIE	TRENCHING	4 IP TO .18 OZ IN GRABS 1.8% ZN IN GRABS
(13)	1944	A LUMSDEN	SHAFT 10'	SHAFT IN RED PORP. DYKE PINK QZ VEIN AT BOTTOM

LEGEND

PROTEROZOIC

ALKALIC INTRUSIVE ROCKS

7 — 7a Coarse-grained syenite, syenite porphyry, pegmatite. 7b Mafic syenite, syenite contaminated by country rocks. 7c Medium-grained syenite, lamprophyres.

INTRUSIVE CONTACT

ARCHEAN

MAFIC INTRUSIVE ROCKS (MATACHEWAN OR NIPISSING)

6 — 6 Diabase.

INTRUSIVE CONTACT

FELSIC INTRUSIVE ROCKS (ALGOMAN)

5 — 5a Granite, porphyritic granite, granodiorite. 5b Granodioritic gneiss. 5c Syenite and trachyte. 5d Mafic syenites and mafic trachyte.

INTRUSIVE CONTACT

EARLY MAFIC AND ULTRAMAFIC ROCKS (HARLEYBURIAN, EARLY ALGOMAN, AND KEEWATIN)

4 — 4a Serpentinite, peridotite. 4b Gabbro, diorite.

INTRUSIVE CONTACT

METASEDIMENTS (TIMISKAMING AND KEEWATIN)

3 — 3a Conglomerate. 3b Quartzose greywacke. 3c Tuff, agglomerate, minor amounts of trachyte agglomerate, breccia.

UNCONFORMITY AND INTERBEDDING

FELSIC METAVOLCANICS (KEEWATIN)

2 — 2a Rhyolite, dacite. 2b Iron formation, silicic tuff, agglomerate. 2c Dacite accanmy, amygdaloidal and spherulitic dacite.

MAFIC METAVOLCANICS AND METASEDIMENTS

1 — 1 Unassimilated mafic volcanics. 1a Basalt, andesite. 1b Chloritic mafic tuff, agglomerate. 1c Altered (bleached, carbonatized) volcanic and sedimentary rocks. 1d Amphibolite, garnet-bearing amphibolite, amphibolite gneiss. 1e Andesite accanmy, amygdaloidal and spherulitic dacite. 1g Biotite-garnet-pyroxene amphibolite.

SYMBOLS

- Glacial striae.
- Area of bedrock outcrop.
- Bedding, top unknown: (inclined, vertical).
- Lava flow: top (arrow) from pillows shape and zoning.
- Schistosity: (horizontal, inclined, vertical).
- Gneissosity, (horizontal, inclined, vertical).
- Geological boundary, observed.
- Geological boundary, position interpreted.
- Fault: (observed, assumed). Last indicates down throw side, arrows indicate horizontal movement.
- Lineament.
- Drag folds with plunge.
- Anticline, syncline, with plunge.
- Drill hole: (vertical, inclined).
- Vein, vein network. Width in inches.
- Shaft: depth in feet.
- Motor road. Provincial highway number encircled where appropriate.
- Other road.
- Trail, portage or winter road.
- Building.

0 1/4 1/2 Mile

0 500 Metres

1 inch = 1/4 mile

FIGURE 4

PROPERTY GEOLOGY AND EXPLORATION

(After OGS Map 2239)

DRILL HOLE LOCATION MAP

CL# L1239349 EBY TWP.

LOT 2 CONS. 5 S½ of S½






Figure - 5

DRILL HOLE SECTION - FACING EAST

EDH-12 EBY TWP., CL # L1239349





1 INCH = 20 FEET
1 : 240

Figure - 6

PRESENT WORK

During the summer of 2005, a program of core drilling was performed with an "X-Ray" type portable core drill. The prospecting target was the area underlaying a swamp where a contact/fault area between mafic volcanic breccia\flows is thought to occur . Though slow and , the good samples and portability of this system is a major advantage.

A - 45° drill hole was put down to a depth of 107 feet to follow up on a - 60° drill hole on the same site. Neither holes passed through the intended target.

RESULTS/CONCLUSIONS

The drill hole encountered a somewhat strained, brecciated andesite to basalt flow with less than 1% fine to 1mm pyrite throughout.

Ministry of Northern Development and Mines
Ontario

Ministère du Développement du Nord et des Mines

Diamond Drilling Log

Journal de forage au diamant

Complete this form related sketch in duplicate.
Remplir en deux exemplaires la présente formule et le croquis annexé

Fill in on every page
Remplir ces cases à chaque page

Hole No. / Forage n° EDH-12 | Page No. / Page n° 1

Drilling Company / Compagnie de forage: ERIC MARION	Collar Elevation / Élévation du collier: 15/16 CORE	Bearing of hole from true North/Position du forage par rapport au nord vrai: 170° AST	Total Footage / Avancement total du forage: 107	Dip of Hole at / Inclinaison du forage au: Collar/collier -45°	Address/Location where core stored / Adresse/endroit où la carotte est stockée: 126 DUNCAN AVE KIRKLAND LAKE ONT.	Map Reference No. / N° de référence sur la carte: —
Date Hole Started / Date de commencement du forage: JULY 4 2005	Date Completed / Date d'achèvement: JULY 26, 2005	Date Logged / Date d'inscription au journal: JAN 15, 2007	Logged by / Inscrit par: E. MARION	Ft./Pi		Claim No. / N° de concession minière: 1239349
Exploration Co., Owner or Optionee / Compagnie d'exploration, propriétaire ou titulaire d'option: ERIC MARION		Date Submitted / Date de dépôt: FEB 19/07	Submitted by (Signature) / Déposé par (signature): [signature]	Ft./Pi		Location (Twp. Lot, Con. or Lat. and Long.) / Emplacement (canton, lot, concession, ou latitude et longitude): EBY TOWNSHIP LOT 2 CONSESSION 5 SE 1/4 of S 1/2
				Ft./Pi		Property Name / Nom de la propriété: PLINKY - WEST

Footage/Avancement From/De	To/À	Rock Type / Type de roche	Description (Colour, grain size, texture, minerals, alteration, etc.) / Description (Couleur, granulométrie, texture, minéraux, transformation, etc.)	Planar Feature Angle*/Angle des caractéristiques planes	Core Specimen Footage † / Longueur en pieds des carottes prélevées	Your Sample No. / N° d'échantillon du prospecteur	Sample Footage/Niveau de prélèvement de l'échantillon (en pieds) From/De	To/À	Sample Length / Longueur de l'échantillon	Assays † / Analyses minéralurgiques
0	107	BASALT - BRECCIA	MEDIUM TO FINE GRAINED GREEN TO DARK GREEN/GREY THIN BASALT FLOWS. BRECIATED OR AGGLOMERATIC SECTIONS BETWEEN FLOWS, ROCK GENERALLY AROUND HARDNESS OF KNIFE WITH SILICEOUS BRECCIA SECTIONS TO HARDNESS OF FILE. LIGHT CARBONATE ALTERATION (10% HCL LIGHT BUBLING) IN SOFTER SECTIONS. LIGHT HEMATITE ALT. IN BRECCIA AS REDDISH/BRN STREAKS AND TINT IN CORE. QZ-CALCITE STRINGERS TO 1/8 THROUGHOUT AT VARIOUS ANGLES. GRAIN OR STRAIN IN BRECCIA SECTIONS DOMINANTLY AT ABOUT 30° TO C.A. NUMEROUS 1/8 STRINGERS OF CALCITE + FLESHY COLORED HEMATITE AT GENERALLY HIGH ANGLES TO CORE. LESS THAN 1% Pyrite THROUGHOUT HOLE BUT UP TO 1% IN BRECCIA SECTIONS. ODD WISPY PYRITE STRINGERS (FINE GRAIN) RANDOMLY THROUGH HOLE IN FLOW + BRECCIA SECTIONS.			C1	46½	48	1½'	BRECCIA, Hem, Ca, Py
						C2	48'	51'	3'	BRECCIA, Hem, Ca, Py
						C3	52½	53½	1'	BRECCIA, Hem, Ca, Py
						C4	77½	80	2½'	BRECCIA, Hem, Ca, Py
						C5	92½	95	2½'	BRECCIA Hem, CaPy

0204 (03/91)

*For features such as foliation, bedding, schistosity, measured from the long axis of the core.
*Exemples de caractéristiques : foliation, schistosité, stratification. L'angle est mesuré par rapport à l'axe longitudinal de la carotte.

† Additional credit available. See Assessment Work Regulation.
† Des crédits supplémentaires sont offerts. Consulter les règlements relatifs aux travaux d'évaluation.
Nota : Dans cette formule, lorsqu'il désigne des personnes, le masculin est utilisé au sens neutre.

Swastika Laboratories Ltd

Established 1928

Assaying - Consulting - Represent...

Geochemical Analysis Certificate

7W...-RG1

Company: **ERIC MARION**
Project: EDH-2
Attn: E. Marion

Date: F...-07-07

We hereby certify the following Geochemical Analysis of 5 Core san... submitted JAN-16-07 by .

Sample Number	Au PPB	Au Check PPB	Multi Element
C-1	Nil	-	Results
C-2	Nil	Nil	to
C-3	7	-	follow
C-4	Nil	-	
C-5	Nil	-	

Certified by Denis Thorne

1 Cameron Ave., P.O. Box 10, Swastika, Ontario P0K 1T0
Telephone (705) 642-3244 Fax (705) 642-3300

Bibliography

Barnes, S.J., Gorton, M.P. and Naldrett, A.J.
1981:Platinum Group Elements in Abitibi Komatiites Associated with Nickle Sulfide Deposits; in Geoscience Research Grant Program, Summary of Research, 1980 to 1981, Ontario Geological Survey,Miscellaneous Paper 98, Grant 17, p. 1-12

Crocket, J.H. and Bowins, R.J.
1985:Rare Earth Element Properties of Archean Iron Formations and their Host Rocks-Some Results from the Temagami and Boston Iron Formations; in Geoscience Research Grant Program, Summary of Research 1984 to 1985, Ontario Geological Survey, Miscellaneous Paper 127, Grant 132, p. 10-14

Crocket,J.H., Blum,N., Hurley, R.,Bowins,R., McRoberts,G., Fyon,A., McNutt,R.H., Schwarcz,H.P. and Rees, C.E.
1984:Geological and Geochemical Studies of the Boston and Temagami Iron Formations and their Contiguous Volcanosedimentary Piles; in Geoscience Research Grant Program, Summary of Research, 1983 to 1984,Ontario Geological Survey,Miscellaneous Paper 121, Grant 132, p. 72-83

Crocket,J.H., Blum,N., Hurley, R.,Bowins,R., McRoberts,G., Fyon,A., McNutt,R.H., Schwarcz,H.P. and Rees, C.E.
1983:Isotopic and Geochemical Characterization of Archean Iron Formations and Associated Volcanic Rocks-Some Preliminary Results from the Temagami and Boston Iron Formations; in Geoscience Research Grant Program, Summary of Research, 1982 to 1983,Ontario Geological Survey,Miscellaneous Paper 113,, Grant 132,p.29-40

Fyon,J.A., Crocket, J.H., Schwarcz, H.P., Kabir, A., and Knyf, M.
1981: Trace Element and Stable Isotope Geochesistry of Auriferous Iron Formations in the Timmins Area; in Geoscience Research Grant Program, Summary of Research, 1980 to 1981, Ontario Geological Survey Miscellaneous Paper 98, Grant 49, p. 90-107

Fortescue, J.A.C. and Gleeson, C.F.
1984: An introduction to the Kirkland Lake (KLIP) Basil Till Geochemical and Mineralogical Study (1979-1982), Timiskaming District, Ontario Geological Survey, Map 80 714, Geochemical Series, Compiled 1984

Hicks,K.D.,and Hattori, K.
1988:Magmatic-Hydrothermal and Wall Rock Alteration Petrology at the Lake Shore Gold Deposit,Kirkland Lake, Ontario; in Geoscience Research Grant Program, Summary of Research 1987 to 1988, Ontario Geological Survey, Miscellaneous Paper 140,Grant 313, p.192-204

Hattori, Keiko, and Levesque, G.
1989: Hydrothermal Activity in the Kirkland Lake Intrusive Complex,Temiskaming District, Ontario, in Geoscience Research Grant Program,Summary of Research 1988 to1989, Ontario Geological Survey, Miscellaneous Paper 143, Grant 313, p. 59-67

Jensen, L.S. and Langford, F.F
1983:Geology and Petrogenesis of the Archean Abitibi Belt in the Kirkland Lake Area, O.G.S. Open File Report 5455

Lovell, H.L.
1972: Geology of the Eby and Otto Area, District of Temiskaming, Ontario Departnen of Mines and Northern Affairs, Geological Report 99, Accompanied by Map 2239, scale 1 inch to 1/2 mile

Lovell, H.L. and Caine, T.W.
1970:Lake Temiskaming Rift Valley; Ontario Department of Mines Miscellaneous Paper 39

Moore, J.C.G.
1966: Geology of Burt Holmes Area,District of Temiskaming,Ontario Department of Mines Geological Report 44, Accompanied by Map2078, scale 1 inch to 1/2 mile

O.G.S.
1979:Airbourne Electromagnetic and Total Intensity Magnetic Survey, Kirkland Lake Area, Morrisette Township, District of Temiskaming : by Questor Surveys Limited for the Ontario Geological Survey, Prelim.Map P.2258 Geophys. Ser., Scale 1:20,000, Survey and compilation February and March 1979

O.G.S.
1979:Airbourne Electromagnetic and Total Intensity Magnetic Survey, Kirkland Lake Area, Teck Township, District of Temiskaming : by Questor Surveys Limited for the Ontario Geological Survey, Prelim.Map P.2263A North Half, and 2263B South Half ,Geophys. Ser., Scale 1:20,000, Survey and compilation February and March 1979

O.G.S.
1986: Volcanology and Mineral Deposits, Miscellaneous Paper 129

Powell, W.G., Hodgson, C.J., and Hanes, J.A.
1989:The Expression of the Larder Lake Break in the Matachewan Area,Temiskaming District,Ontario; in Geoscience Research Grant Program, Summary of Research 1988 to 1989,Ontario Geological Survey, Miscellaneous Paper 143, Grant 329, p. 125-132

Savage, W.S.
1964:Mineral Resources and Mining Properties in the Kirkland Lake - Larder Lake Area, District of Temiskaming, Ontario Department of Mines Mineral Resource Circular No. 3 Accompanied by Chart A Kirkland-Larder Lake Area, scale 1 inch to 2 miles

Toogood, D.J. and Hodgson, C.J.
1986: Relationship Between Gold Deposits and the Tectonic Framework of the Abitibi Greenstone Belt in the Kirkland Lake-Larder Lake Area; in Geoscience Research Grant Program, Summary of Research 1985 to 1986,Ontario Geological Survey, Miscellaneous Paper 130,Grant 227, p.79-86

Ministry of Northern Developement and Mines
:Resident Geologist Files, literally hundreds of files of submitted assesment work files covering many claim areas in Bernhardt Township, Eby Township, Grenfell Township, Gauthier Township, Lebel Township, Morrisette Township, Otto Township, Teck Township,

O.G.S.
1986: Volcanology and Mineral Deposits, Miscellaneous Paper 129